

Initial Date: 11/15/2012 Revised Date: 05/25/2023

Michigan ADULT CARDIAC BRADYCARDIA

Bradycardia

-∿-

This protocol is for paramedic use only

This is a protocol for patients with serious symptomatic bradycardia, defined as patients with heart rate less than 60 bpm and hypotension, or shock. Titrate treatments to a heart rate above 60 bpm. If the patient remains hypotensive, refer to the **Shock Treatment Protocol**.

- 1. Follow the General Pre-Hospital Care-Treatment Protocol.
- 2. Administer **atropine** 1 mg IV/IO rapid push repeating every 3-5 minutes to a total dose of 3 mg IV/IO, until a heart rate of greater than 60/minute is reached.
- 3. Transcutaneous pacing (TCP) when available may be initiated prior to establishment of IV access and/or before **atropine** begins to take effect. Pacing is the treatment of choice for high degree A-V block (second-degree Type II, or third-degree), apply pacer pads. Follow the **Electrical Therapy- Procedure Protocol.**
- 4. Per MCA selection, provide sedation per **Patient Procedural Sedation-Procedure Protocol**
- 5. For patients with persistent symptomatic bradycardia, administer **epinephrine** by push dose (dilute boluses)
 - a. Prepare (10 mcg/mL) by adding 1mL of 1mg/10mL **epinephrine** in 9mL **NS**, then:
 - i. Administer 10-20 mcg (1-2 mL epinephrine 10 mcg/mL) IV/IO
 - ii. Repeat every 3 to 5 minutes
 - iii. Titrate SBP greater than 90 mmHg

Notes:

- 1. Consider possible etiologies:
 - A. Hyper/hypokalemia, other metabolic disorders
 - B. Hypothermia
 - C. Hypovolemia (including vomiting/diarrhea)
 - D. Hypoxia
 - E. Hydrogen ion excess (acidosis)
 - F. Toxins/ overdose (e.g., beta-blocker or calcium channel-blocker)
 - G. Tamponade
 - H. Tension pneumothorax
 - I. Thrombosis (pulmonary or coronary)
- 2. Transcutaneous pacemaker electrode pads may be applied to these patients without initiating pacing so that the pacemaker is ready if patient condition deteriorates.
- 3. For symptomatic high-degree (second-degree Type II, or third-degree) AV block, begin pacing without delay.
- 4. Heart transplant patients may not respond to **atropine**

Medication Protocols Atropine Epinephrine

Protocol Source/References: Highlights of the 2020 AHA Guidelines Update for CPR and ECC