

Pulmonary Edema/Cardiogenic Shock

This protocol is to be followed for patients in respiratory distress due to pulmonary edema with or without hypotension (i.e., CHF/fluid overload or Cardiogenic Shock). Pulmonary edema usually presents with crackles which should be continuously evaluated as they may evolve with treatments.

1. Follow **General Pre-Hospital Care-Treatment Protocol**.
2. Initiate supplemental oxygen by non-rebreather mask.
3. Position patient upright with legs dependent, if possible.
4. Consider CPAP per **CPAP-Procedure Protocol**
5. Establish IV access without delaying treatment per **Vascular Access & IV Fluid Therapy-Procedure Protocol**.
6. If wheezing, administer **albuterol** 2.5 mg/3ml **NS** nebulized (Per MCA selection may be EMT skill) per **Medication Administration-Medication Protocol**

Nebulized **albuterol** administration per
MCA selection
☐ EMT

7. If crackles (with or without wheezing) administer **nitroglycerin** as outlined below.
 - a. Inquire of all patients regardless of identified gender if they have taken an erectile dysfunction medication or medications used to treat pulmonary hypertension in the last 48 hours.
 - i. If yes, DO NOT ADMINISTER NITROGLYCERIN AND CONTACT MEDICAL CONTROL.
 - b. Prior to IV administration if no erectile dysfunction medication and systolic BP is above 120 mmHG, **nitroglycerin** 0.4mg sublingual may be administered up to a maximum of 3 doses.
 - c. If SBP above 100 mmHg (with IV/IO in place), administer **nitroglycerin** 0.4 mg SL, repeat every 3-5 minutes if SBP remains above 100 mmHg.
 - d. If wheezing continues, continue **nitroglycerin** 0.4 mg SL and consider: **albuterol/ipratropium bromide** per **Respiratory Distress-Treatment Protocol**
8. If SBP is below 100 mmHG treat for cardiogenic shock.
 - a. Prepare (**epinephrine** 10 mcg/mL) by combining 1mL of 1mg/10mL **epinephrine** in 9mL **NS**
 - i. Administer 20 mcg (2 mL **epinephrine** 10 mcg/mL) IV/IO
 - ii. Repeat every 3-5 minutes
 - iii. Titrate SBP greater than 90 mm/Hg.
9. If indicated, consider an advanced airway see **Airway Management-Procedure Protocol**.
10. Obtain 12-lead ECG (May be a BLS or Specialist skill, per MCA selection, see **12 Lead ECG-Procedure Protocol**). Follow MCA transport protocol if ECG is positive for ST segment elevation myocardial infarction (STEMI) and alert hospital as soon as possible.

Initial Date: 11/15/2012

Revised Date: 06/03/2023

Section 5-4

Medication Protocols

Albuterol

Epinephrine

Nitroglycerin