

MEDICATION SECTION
FENTANYL

Initial Date: 07/19/2023

Revised Date:

Fentanyl

Pharmacological Category: Analgesic, Opioid; General Anesthetic

Routes: IV/IO/IM/IN

Indications:

1. Pain management
2. Patient sedation

Contraindications:

1. Altered Mental Status
2. Hypotension
3. Respiratory Depression

Expected effects:

1. Decreased pain
2. Decreased agitation

Side effects:

1. Drowsiness
2. Hypotension
3. Respiratory Depression
4. Vomiting

Dosing: CHEST PAIN/ACUTE CORONARY SYNDROME

Indication: Chest pain in which nitroglycerin is contraindicated due to erectile dysfunction medication or suspected cardiac chest pain is refractory to nitroglycerin.

Adults (65 years of age or under) administer:

1. Fentanyl 1 mcg/kg IV/IO/IN, max single dose 100 mcg. May repeat one time.
Total dose may not exceed 200 mcg.

Adults (> 65 years of age or older) administer:

1. Fentanyl 0.5 mcg/kg IV/IO/IN. Max single dose 50 mcg. May repeat three times.
Total dose may not exceed 200 mcg.

Dosing: PAIN MANAGEMENT

Indication: Patient is unable to tolerate ketamine or ketamine is not available and the patient has significant pain (described as 7 or greater on the Wong Pain Scale).

Adults 65 years of age or under administer:

1. Fentanyl 1 mcg/kg IV/IO/IN. Max single dose 100 mcg. May repeat one time. Total dose may not exceed 200 mcg.

Adults > 65 years of age administer:

MEDICATION SECTION
FENTANYL

Initial Date: 07/19/2023

Revised Date:

1. Fentanyl 0.5 mcg/kg IV/IO/IN. Max single dose 50 mcg. May repeat three times. Total dose may not exceed 200 mcg.

Pediatrics administer:

1. Fentanyl according to MI MEDIC cards
2. If MI MEDIC cards are not available administer:
 - a. Fentanyl 0.1 mg/kg IV/IO/IN

Dosing: PATIENT PROCEDURAL SEDATION

Adults administer:

1. Fentanyl 50-100 mcg (1 mcg/kg) IV/IO titrated slowly (IN, if available). May repeat every 4 minutes to a maximum of 3 mcg/kg.

Pediatrics administer:

1. Fentanyl according to MI MEDIC cards
2. If MI MEDIC cards are not available administer:
 - a. Fentanyl 1 mcg/kg IV/IO titrated slowly (IN, if available). May repeat every 5 minutes to a maximum of 3 mcg/kg.

Used in the Following Protocols

Chest Pain/Acute Coronary Syndrome (Section 5 Adult Cardiac)
Pain Management (Section 7 Procedures)
Patient Procedure Sedation (Section 7 Procedures)