

Initial Date: 11/15/2012
Revised Date: 03/03/2025

Section 3-4

Seizures

For patients $\leq$ 14 years of age refer to **Pediatric Seizure-Treatment Protocol**

1. Follow **General Pre-hospital Care-Treatment Protocol**.
2. IF PATIENT IS ACTIVELY SEIZING:
 - A. Protect patient from injury.
 - B. Do not force anything between teeth.
 - C. Pregnant women 20 weeks gestation up to 6 weeks post birth WITHOUT a seizure disorder history, treat as eclampsia, see **magnesium sulfate** administration below (C.)
 -  D. Administer **midazolam** 10 mg IM prior to IV start
-  3. Check blood glucose (may be MFR skill, see **Blood Glucose Testing-Procedure Protocol**),
 -  A. If blood glucose is found to be less than 60 mg/dL or hypoglycemia is suspected administer **dextrose** 25 gm IV per **Dextrose-Medication Protocol**
 -  B. If no IV access, per MCA selection, administer **glucagon** 1 mg IM or 3 mg IN (if available per MCA selection), (may be EMT skill per MCA selection).

Glucagon administration per MCA Selection

☐ Not included

	1 mg glucagon IM *Injectable formulation ONLY*	3 mg glucagon IN *Intranasal formulation ONLY*
EMT	☐	☐
Specialist	☐	☐
Paramedic	☐	☐

*INTRANASAL GLUCAGON ADMINISTRATION: Only glucagon that is FDA-approved for nasal administration (e.g., Baqsimi(R)) may be given by IN route. Injectable glucagon is not to be administered via IN route. EMS clinicians may assist family/patient care givers in administering glucagon that is FDA-approved for IN use, if prescribed for the patient (regardless of age).

If patient is pregnant (eclampsia)

-  a. Administer **magnesium sulfate** 4 gm over 10 minutes IV/IO until seizure stops. Administration of **magnesium sulfate** is best accomplished by adding **magnesium sulfate** 4 gm to 100 or 250 ml of **NS** and infusing over approximately 10 minutes.
- b. If eclamptic seizure does not stop after magnesium, then administer benzodiazepine as specified below.
-  C. If IV already established and **midazolam** IM/IN has not been administered, administer **midazolam** 5 mg IV/IO

Initial Date: 11/15/2012

Revised Date: 03/03/2025

Section 3-4



D. If seizures persist



a. Repeat **midazolam** 5mg IV/IO/IM/IN

b. Contact Medical Control

4. IF PATIENT IS NOT ACTIVELY SEIZING and has/is:

A. Altered level of consciousness, refer to **Altered Mental Status-Treatment Protocol**.

B. Alert



a. Monitor for changes.

b. Obtain vascular access.



c. Check blood glucose (may be MFR skill, see **Blood Glucose Testing-Procedure Protocol**),

Medication Protocols

Dextrose

Glucagon

Magnesium Sulfate

Midazolam

Protocol Source/References: NAEMSO Clinical Guidelines